




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : A61K 39/39 // (A61K 39/39, 47:40, 47:34)		(11) International Publication Number: WO 99/10008
A1		(43) International Publication Date: 4 March 1999 (04.03.99)
(21) International Application Number: PCT/US98/1794C		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GH, GM, HR, HU, ID, IL, IS, JP, KB, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KR, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BB, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
(22) International Filing Date: 28 August 1998 (28.08.98)		
(30) Priority Data: 60/057,255 29 August 1997 (29.08.97) US		
(71) Applicant: AQUILA BIOPHARMACEUTICALS, INC. [US/US]; 365 Plantation Street, Worcester, MA 01605-2376 (US).		
(72) Inventors: KENSIL, Charlotte; 15 Camp Street, Milford, MA 01757 (US). BELTZ, Gerald, A.; 4 Eustis Street, Lexington, MA 02173 (US).		
(74) Agents: SUPERKO, Colleen et al.; Hale and Dorr LLP, 60 State Street, Boston, MA 02109 (US).		Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>

(54) Title: COMPOSITIONS COMPRISING THE ADJUVANT QS-21 AND POLYSORBATE OR CYCLODEXTRIN AS EXCIPIENT



(57) Abstract

Certain novel compositions of the adjuvant saponin QS-21 having improved properties are disclosed. The compositions of the present invention are designed (1) to minimize the lytic effects of QS-21, (2) to improve the tolerance of QS-21 containing formulations in humans or other animals, (3) to stabilize the QS-21 from alkaline hydrolysis and/or (4) to maintain the high adjuvant potency of the QS-21 product. These compositions may be employed with vaccines comprising proteins or peptides, polysaccharides, lipids, or nucleic acids.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SL	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	ME	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

COMPOSITIONS COMPRISING THE ADJUVANT QS-21 AND POLYSORBATE OR CYCLODEXTRIN
AS EXIPIENTFIELD OF THE INVENTION

5 The present invention relates to the field of immune adjuvants and the
use thereof as immune adjuvants in vaccines. The compositions of the present
invention exhibit significantly improved properties relevant to the lytic effect,
tolerance to QS-21 associated pain, and product stability of QS-21, and
10 maintain full adjuvant activity.

BACKGROUND OF THE INVENTION

15 Adjuvant saponins have been identified and purified from an aqueous
extract of the bark of the South American tree, *Quillaja saponaria* Molina.
Among the 22 peaks which were separable and displayed saponin activity,
QS-21 was one of the more predominant purified saponins. This saponin has
20 been substantially purified by high pressure liquid chromatography (HPLC),
low pressure liquid silica chromatography, and hydrophilic interactive
chromatography (HILIC). QS-21 has been found to be useful as an immune
adjuvant for enhancing immune responses in individuals at a much lower
25 concentration than the previously available heterogeneous saponin preparations
without the toxic effects associated with crude saponin preparations.

 QS-21 is a membrane-lytic triterpene glycoside saponin. It forms
micelles of approximately the same radius as bovine serum albumin (Kensil,
30 U.S. Patent No. 5,057,540) and has a critical micellar concentration of
approximately 50 µg/ml in PBS (Soltysik, S., et al., 1995, Vaccine 13:1403-1410).

The potency of an adjuvant formulation containing an antigen plus QS-21 can be assessed in experiments that address the relationship of adjuvant dose to immunological function (dose-response experiments). A decrease in adjuvant potency is expected to increase the minimum dose (threshold dose) required for enhancement of immune response. A desirable composition is expected to maintain an equivalent or better potency than the formulation that is used as a reference. For QS-21, the reference formulation is a simple solution in phosphate-buffered saline (PBS) or saline.

The adjuvant activity of QS-21 is assessed in animal models such as mice. The primary responses measured are increases in antigen-specific antibody and antigen-specific cytotoxic T lymphocytes (CTL). The threshold dose of QS-21 that will enhance murine immune response (antibody or CTL) has been measured in simple buffer solution such as PBS. A dose of 2.5 µg has been shown to be the threshold dose for antibody (Kensil, C.R., et al., 1993, Vaccine Research 2:273-281) and for CTL (Newman, M.J., et al., 1992, Immunology 148:2357-2362) to the antigen ovalbumin (OVA) in C57BL/6 mice in PBS. Similar threshold doses were observed when aluminum hydroxide was included in the PBS formulation (Kensil, C.R., et al., 1993, Vaccine Research 2:273-281). However, it is expected that there may be differences in potency between different compositions of a given adjuvant.

Despite these beneficial qualities, QS-21 possesses some unwelcome qualities as well. For instance, QS-21 associates with phospholipid bilayers and causes a lytic effect on certain cell membranes (i.e., erythrocytes). QS-21 will absorb to the phospholipid bilayer of sheep erythrocytes and cause the red blood cells to release hemoglobin. This hemoglobin release, which is known as

hemolysis, occurs at a concentration of approximately 5-7 µg/ml in a simple buffer such as saline or PBS (Kensil, C.R., et al., 1991, J. Immunology 146:431-437). At higher concentrations (above the critical micellar concentration of QS-21), total lysis of the red blood cell membrane occurs. The lytic effect of QS-21 is, therefore, an undesirable property for a composition.

In *in vivo* studies, hemolysis is not noted. However, after intramuscular injection of QS-21/saline solutions into New Zealand white rabbits, mild to moderate fibroblast damage or necrosis is noted in some animals when the injection site is analyzed histopathologically (Kensil, C.R., et al., 1995, In: Vaccine Design: The Subunit and Adjuvant Approach, Powell, M.F. and Newman, M.J., Eds., Plenum Press, NY). Further, creatine kinase, a marker for muscle damage is increased after injection with QS-21 in saline or PBS. This rise is believed to be due to the lytic effect of QS-21 on cell membranes.

Moreover, in clinical trials, some individuals have experienced an immediate, transient pain after injection with QS-21 in simple buffer solutions (saline or PBS). This pain, described by most individuals as a burning pain, may be a secondary reaction correlated with the lytic effect of the QS-21 adjuvant. Patient pain is likewise an objectionable property for a composition.

Product stability is another concern for QS-21 containing compositions. The shelf life of a vaccine product is typically defined by the extent of time to reach a defined and acceptable low level of degradation (such as, the time to 10% degradation, also known as t_{90}). Most commercial vaccine products have a shelf life of at least 18 to 24 months when stored in the refrigerator at 4°C.

Adjuvants, which are essential components of vaccines, therefore must also have equally long shelf lives. However, the shelf life of a 50 µg/ml solution of

QS-21 at pH 7.0 at 4°C is reached in about 3 months. The reason for the short shelf life is because the ester bond of QS-21 is increasingly labile at increasing pH and because monomers of QS-21, as opposed to micelles, are subject to hydrolysis. The need to stabilize compositions of QS-21 adjuvant is significant.

SUMMARY OF THE INVENTION

A need exists for compositions of the saponin adjuvant QS-21 that may be used to boost the antigenic immune response in a relatively low dose with low local reactions and side effects, but also features a reduced lytic effect, improved tolerance to QS-21, and an increased stability. Accordingly, the present invention provides novel compositions of QS-21 that have these improved characteristics compared to a simple solution of QS-21 in a buffer such as saline or PBS. Surprisingly, the full adjuvant potency of QS-21 in the disclosed compositions is not compromised compared to a control formulation of QS-21 in PBS.

DESCRIPTION OF THE FIGURES

Figure 1 depicts a graph showing the adjuvant potency of various compositions. Figure 1A shows the effect of Polysorbate 40, Polysorbate 60, and Polysorbate 80 on the immune response of Balb/c mice to ovalbumin at different concentrations of QS-21. Figure 1B shows the effect of methyl- β -cyclodextrin on the immune response of Balb/c mice to ovalbumin at different concentrations of QS-21.

Figure 2 depicts a graph showing the effect of Polysorbate 80 and hydroxypropyl- β -cyclodextrin on Type 14 IgG3 antibody response to a

T-independent polysaccharide antigen.

Figure 3 shows a bar graph of patients' tolerance to pain for various excipients in QS-21 adjuvant compositions from Trial 1. This figure shows how the pain scores are classified as no pain, mild pain, moderate pain, or severe pain, where 0=no pain, 1-3=mild pain, 4-7=moderate pain, and 8-10=severe pain.

Figure 4 shows the individual scores for the patients' tolerance to pain in Figure 3. This figure shows individual immediate pain scores after injection of a given formulation on a scale of 0-10, where 0 is no pain and 10 is maximum pain.

Figure 5 shows a bar graph of patients' tolerance to pain for various excipients in QS-21 adjuvant compositions from Trial 2. This figure shows how the pain scores are classified as no pain, mild pain, moderate pain, or severe pain, where 0=no pain, 1-3=mild pain, 4-7=moderate pain, and 8-10=severe pain.

Figure 6 shows the individual scores for the patients' tolerance to pain in Figure 5. This figure shows individual immediate pain scores after injection of a given formulation on a scale of 0-10, where 0 is no pain and 10 is maximum pain. Mean and median scores for each formulation are listed below each formulation.

DESCRIPTION OF THE PREFERRED EMBODIMENTS

The saponins of the present invention may be obtained from the tree *Quillaja saponaria* Molina.

The term "saponin" as used herein includes glycosidic triterpenoid

compounds which produce foam in aqueous solution, have hemolytic activity in most cases, and possess immune adjuvant activity. The invention encompasses the saponin per se, as well as biologically active fragments thereof.

The invention also concerns compositions, such as immunologic compositions, comprising one or more substantially pure saponin fractions, and methods of using these compositions as immune adjuvants.

More particularly, the compositions of the present invention may reduce the *in vitro* lytic effects of a saponin adjuvant containing formulation. Another preferred composition is one that may maintain the maximum adjuvant activity of a saponin. Yet another preferred composition may increase the stability of a saponin adjuvant containing composition from alkaline hydrolysis. Other compositions may preferably improve an individual's tolerance to saponin adjuvant associated pain from a formulation containing a saponin adjuvant.

As described in Kensil, et al., U.S. Patent No. 5,057,540, the contents of which are fully incorporated by reference herein, the adjuvant activity of such saponins may be determined by any of a number of methods known to those of ordinary skill in the art. The increase in antibody titer of antibody against specific antigen upon administration of an adjuvant may be used as a criteria for adjuvant activity. (Dalsgaard, *Acta Veterinaria Scandinavica*, 69:1 (1978); Bomford, *Int. Archs. Allergy Appl. Immun.* 77:409 (1985).) Briefly, one such test involves injecting CD-1 mice intradermally with an antigen (for instance, i.e., bovine serum albumin, BSA) mixed with varying amounts of the potential adjuvant. Sera was harvested from the mice two weeks later and tested by ELISA for anti-BSA antibody.

"QS-21" designates the mixture of isomeric components QS-21-V1 and QS-21-V2 which appear as a single peak on reverse phase HPLC on Vydac C4 (5 μ m particle size, 300Å pore, 4.6 mm ID x 25 cm) in 40 mM acetic acid in methanol/water (58/42, v/v). The component fractions are referred to specifically as QS-21-V1 and QS-21-V2 when describing experiments performed on the further purified components.

The term "substantially pure" means substantially free from compounds normally associated with the saponin in its natural state and exhibiting constant and reproducible chromatographic response, elution profiles, and biologic activity. The term "substantially pure" is not meant to exclude artificial or synthetic mixtures of the saponin with other compounds.

The substantially pure QS-7 saponin also referred to as QA-7 in U.S. Patent No. 5,057,540) is characterized as having immune adjuvant activity, containing about 35% carbohydrate (as assayed by anthrone) per dry weight, having a UV absorption maxima of 205-210 nm, a retention time of approximately 9-10 minutes on RP-HPLC on a Vydac C₄ column having a 5 μ m particle size, 300 Å pore, 4.6 mm ID x 25 cm L in a solvent of 40 mM acetic acid in methanol/water (58/42; v/v) at a flow rate of 1 ml/min, eluting with 52-53% methanol from a Vydac C₄ column having a 5 μ m particle size, 300 Å pore, 10 mm ID x 25 cm L in a solvent of 40 mM acetic acid with gradient elution from 50 to 80% methanol, having a critical micellar concentration of approximately 0.06% (w/v) in water and 0.07% (w/v) in phosphate buffered saline, causing no detectable hemolysis of sheep red blood cells at concentrations of 200 μ g/ml or less, and containing the monosaccharide residues terminal rhamnose, terminal xylose, terminal glucose, terminal

galactose, 3-xylose, 3,4-rhamnose, 2, 3-fucose, and 2,3-glucuronic acid, and apiose (linkage not determined).

The substantially pure QS-17 saponin (also referred to as QA-17 in U.S. Patent N. 5, 057, 540) is characterized as having immune adjuvant activity, containing about 29% carbohydrate (as assayed by anthrone) per dry weight, having a UV absorption maxima of 205-210 nm, a retention time of approximately 35 minutes on RP-HPLC on a Vydac C₄ column having a 5 µm particle size, 300 Å pore, 4.6 mm ID x 25 cm L in a solvent of 40 mM acetic acid in methanol/water (58/42; v/v) at a flow rate of 1 ml/min, eluting with 63-64% methanol from a Vydac C₄ column having a 5 µm particle size, 300 Å pore, 10 mM ID x 25 cm L in a solvent of 40 mM acetic acid with gradient elution from 50 to 80% methanol, having a critical micellar concentration of approximately 0.06% (w/v) in water and 0.03% (w/v) in phosphate buffered saline, causing hemolysis of sheep red blood cells at 25 µg/ml or greater, and containing the monosaccharide residues terminal rhamnose, terminal xylose, 2-fucose, is characterized as having immune adjuvant activity, containing about 35% carbohydrate (as assayed by anthrone) per dry weight, having a UV absorption maxima of 205-210 nm, a retention time of approximately 9-10 minutes on RP-HPLC on a Vydac C₄ column having a 5 µm particle size, 300 Å pore, 4.6 mm ID x 25 cm L in a solvent of 40 mM acetic acid in methanol/water (58/42; v/v) at a flow rate of 1 ml/min, eluting with 52-53% methanol from a Vydac C₄ column having a 5 µm particle size, 300 Å pore, 10 mM ID x 25 cm L in a solvent of 40 mM acetic acid with gradient elution from 50 to 80% methanol, having a critical micellar concentration of approximately 0.06% in water and 0.07% in phosphate buffered saline, causing no detectable

hemolysis of sheep red blood cells at concentrations of 200 µg/ml or less, and containing the monosaccharide residues terminal rhamnose, terminal xylose, 2-fucose, 3-xylose, 3,4-rhamnose, 2,3-glucuronic acid, terminal glucose, 2-arabinose, terminal galactose and apiose (linkage not determined).

The substantially pure QS-18 saponin (also referred to as QA-18 in U.S. Patent No. 5,057,540) is characterized as having immune adjuvant activity, containing about 25-26% carbohydrate (as assayed by anthrone) per dry weight, having a UV absorption maxima of 205-210 nm, a retention time of approximately 38 minutes on RP-HPLC on a Vydac C₄ column having a 5 µm particle size, 300 Å pore, 4.6 mm ID x 25 cm L in a solvent of 40 mM acetic acid in methanol/water (58/42; v/v) at a flow rate of 1 ml/min, eluting with 64-65% methanol from a Vydac C₄ column having a 5 µm particle size, 300 Å pore, 10 mM ID x 25 cm L in a solvent of 40 mM acetic acid with gradient elution from 50 to 80% methanol, having a critical micellar concentration of approximately 0.04% (w/v) in water and 0.02% (w/v) in phosphate buffered saline, causing hemolysis of sheep red blood cells at 25 µg/ml or greater, and containing the monosaccharide residues terminal arabinose, terminal apiose, terminal xylose, terminal glucose, terminal galactose, 2-fucose, 3-xylose, 3,4-rhamnose, and 2,3-glucuronic acid.

The substantially pure QS-21 saponin (also referred to as QA-21 in U.S. Patent No. 5,057,540) is characterized as having immune adjuvant activity, containing about 22% carbohydrate (as assayed by anthrone) per dry weight, having a UV absorption maxima of 205-210 nm, a retention time of approximately 51 minutes on RP-HPLC on a Vydac C₄ column having a 5 µm particle size, 300 Å pore, 4.6 mm ID x 25 cm L in a solvent of 40 mM acetic acid

in methanol/water (58/42; v/v) at a flow rate of 1 ml/min, eluting with 69-70% methanol from a Vydac C₁₈ column having a 5 µm particle size, 300 Å pore, 10 mM ID x 25 cm L in a solvent of 40 mM acetic acid with gradient elution

5 from 50 to 80% methanol, having a critical micellar concentration of approximately 0.03% (w/v) in water and 0.02% (w/v) in phosphate buffered saline, causing hemolysis of sheep red blood cells at 25 µg/ml or greater. The component fractions, substantially pure QS-21-V1 and QS-21-V2 saponins, have

10 the same molecular weight and identical spectrums by FAB-MS. They differ only in that QS-21-V1 has a terminal apiose which is xylose in QS-21-V2 (which therefore has two terminal xyloses and no apiose). The two components additionally contain the monosaccharides terminal arabinose, terminal apiose,

15 terminal xylose, 4-rhamnose, terminal galactose, 2-fucose, 3-xylose, and 2,3-glucuronic acid.

The invention may also encompass impure forms of saponin adjuvants.

20 For example, one preferred embodiment is the heterogenic saponin adjuvant known as "Quil A." Commercial preparations of Quil A are available from Superfos (Vedbaek, Denmark) and have been isolated from the bark of the South American tree, *Quillaja saponaria* Molina. Quil A is characterized

25 chemically as carbohydrate moieties in glycosidic linkage to the triterpenoid quillaic acid. Quil A possesses immune adjuvant activity and separates into 20 discrete peaks by RP-HPLC on Vydac C₁₈ column having a 5 µm particle size, 300 Å pore, 4.6 mM ID x 25 cm L in a solvent of 40 mM acetic acid in methanol

30 water (U.S. Patent No. 5,057,540).

The invention also relates to a composition which comprises a saponin adjuvant of the present invention, an antigen, and an excipient. Preferably, the

adjuvant is QS-21. Preferably, the excipients may be nonionic surfactants, polyvinylpyrrolidone, human serum albumin, aluminum hydroxide, agents with anesthetic action, and various unmodified and derivatized cyclodextrins.

5 More preferably, the nonionic surfactants may include Polysorbate 20, Polysorbate 40, Polysorbate 60, and Polysorbate 80. The polyvinylpyrrolidone may preferably be Plasdane C15, a pharmaceutical grade of polyvinylpyrrolidone. The agent having anesthetic action preferably is benzyl
10 alcohol. A preferred cyclodextrin is a hydroxypropyl- β -cyclodextrin, which reduces QS-21 lysis of red blood cells *in vitro*.

The term "immune adjuvant," as used herein, refers to compounds which, when administered to an individual or tested *in vitro*, increase the
15 immune response to an antigen in the individual or test system to which said antigen is administered. Preferably, such individuals are humans, however, the invention is not intended to be so limiting. Any animal that may
20 experience the beneficial effects of the vaccines of the invention are within the scope of animals which may be treated according to the claimed invention. Some antigens are weakly immunogenic when administered alone or are toxic to the individual at concentrations which evoke immune responses in said
25 individual. An immune adjuvant may enhance the immune response of the individual to the antigen by making the antigen more strongly immunogenic. The adjuvant effect may also lower the dose of said antigen necessary to achieve an immune response in said individual.

30 The saponins of the present invention may be utilized to enhance the immune response to any antigen. Typical antigens suitable for the immune-response provoking compositions of the present invention include antigens

derived from any of the following: viruses, such as influenza, feline leukemia virus, feline immunodeficiency virus, HIV-1, HIV-2, rabies, measles, hepatitis B, or hoof and mouth disease, bacteria, such as anthrax, diphtheria, Lyme disease
5 or tuberculosis; or protozoans, such as *Babesiosis bovis* or Plasmodium. The antigens may be proteins, peptides, polysaccharides, lipids, or nucleic acids encoding the protein or peptide. The proteins, peptides, lipids, or nucleic acids may be purified from a natural source, synthesized by means of solid phase
10 synthesis, or may be obtained means of recombinant genetics.

Administration of the compounds useful in the method of the present invention may be by parenteral, intravenous, intramuscular, subcutaneous, intranasal, oral or any other suitable means. The dosage administered may be
15 dependent upon the age, weight, species, kind of concurrent treatment, if any, route of administration, and nature of the antigen administered. In general, the saponin and antigen may be administered at a dosage of about 0.001 to about 1.0 mg/kg of saponin adjuvant or antigen per weight of the individual.
20 The initial dose may be followed up with a booster dosage after a period of about four weeks to enhance the immunogenic response. Further booster dosages may also be administered.

25 The effective compound useful in the method of the present invention may be employed in such forms as capsules, liquid solutions, suspensions or elixirs for oral administration, or sterile liquid forms such as solutions or suspensions. The vaccine of the present invention may be administered
30 parenterally, intranasally, or orally.

Another preferred embodiment is a method for reducing the *in vitro* lytic effect of an immune adjuvant composition comprising administering to an

individual an effective amount of QS-21 and an excipient. Preferably, the excipients may be nonionic surfactants, polyvinylpyrrolidone, human serum albumin, aluminum hydroxide, agents with anesthetic action, and various
5 unmodified and derivatized cyclodextrins. More preferably, the nonionic surfactants may include Polysorbate 20, Polysorbate 40, Polysorbate 60, and Polysorbate 80. The polyvinylpyrrolidone may preferably be Plasdone C15, a pharmaceutical grade of polyvinylpyrrolidone. The agent having anesthetic
10 action preferably is benzyl alcohol. A preferred cyclodextrin is Encapsin, a hydroxypropyl- β -cyclodextrin, which reduces QS-21 lysis of red blood cells *in vitro*.

Other preferred methods falling within the scope of the invention
15 include a method for maintaining the maximum adjuvant activity of QS-21 comprising administering to an individual an effective amount of QS-21 and an excipient and a method for improving the tolerance to saponin adjuvant
20 associated pain in an individual to whom it is administered comprising administering an effective amount of QS-21 and an excipient.

EXAMPLES

25 A variety of excipients were evaluated in combination with QS-21 as novel compositions. These included various nonionic surfactants (Triton X-100, Polysorbate 20, Polysorbate 40, Polysorbate 60, and Polysorbate 80), polyvinylpyrrolidone (Plasdone C15), human serum albumin, aluminum
30 hydroxide, agents with anesthetic action (benzyl alcohol), and various unmodified and derivatized cyclodextrins (hydroxypropyl- β -cyclodextrin, hydroxypropyl- γ -cyclodextrin, methyl- β -cyclodextrin). The final formulations

were assessed for their capacity to reduce the lytic effect of QS-21, to improve tolerance to QS-21 adjuvant associated pain in humans, to stabilize QS-21 in aqueous solution, and/or to maintain maximum adjuvant potency relative to a control formulation of QS-21 in PBS.

Example 1

Compositions that Reduce the Lytic Effect of QS-21

A simple *in vitro* assay was used to screen excipients for reducing the lytic effect of QS-21. The lytic effect of QS-21 can be determined in an assay of hemolysis of sheep erythrocytes. Briefly, various two fold serial dilutions of QS-21 in a given excipient are prepared in a round bottom microtiter plate (100 μ l/well). All plates contain control wells containing excipient, but no QS-21. The concentration of QS-21 ranges from 1.56 to 200 μ g/ml. A total volume of 25 μ l of sheep erythrocytes (washed with PBS) is added to each well, mixed with the QS-21/excipient solution, and incubated at ambient temperature for 30 minutes. After the end of the incubation, the round bottom plate is centrifuged at 2000 rpm for 5 minutes to sediment any unlysed cells. A total volume of 75 μ l of supernatant (containing released hemoglobin) is transferred to the equivalent well of a flat-bottom 96 well plate. The flat-bottom plate is centrifuged at 2000 rpm for 5 minutes to break any air bubbles. The absorbance at 570 nm is read in a microtiter plate reader. Absorbance at 570 nm is plotted on the y-axis against QS-21 concentration plotted on the x-axis. The absorbance of hemoglobin in the supernatant of a well where no intact cell pellet was observed is defined as maximum hemolysis. The hemolytic index of QS-21 is defined as the concentration of QS-21 that yields

an absorbance equivalent to 50% of the maximum absorbance. An excipient that reduces the lytic effect of QS-21 is expected to increase the hemolytic index.

5 Table 1 lists the hemolytic indices of QS-21 in various excipients. All excipients were tested in the absence of QS-21. In the absence of QS-21, no hemolysis was noted, indicating that the excipient formulations were isotonic. Excipients that were shown to be effective in minimizing the lytic effect
10 (increase hemolytic index) of QS-21 were hydroxypropyl- β -cyclodextrin, aluminum hydroxide, and Polysorbate 80 in saline.

15

20

25

30

Table 1:

	Excipient	Hemolytic Index (µg/ml)
	PBS	5
5	α-cyclodextrin (2 mg/ml)	1.5
	β-cyclodextrin (2 mg/ml)	10
	methyl-β-cyclodextrin (2 mg/ml)	36
	hydroxypropyl-γ-cyclodextrin (2 mg/ml)	5
	hydroxypropyl-β-cyclodextrin (1 mg/ml)	9
10	hydroxypropyl-β-cyclodextrin (2 mg/ml)	11
	hydroxypropyl-β-cyclodextrin (4 mg/ml)	18
	hydroxypropyl-β-cyclodextrin (8 mg/ml)	32
	hydroxypropyl-β-cyclodextrin (16 mg/ml)	51
15	hydroxypropyl-β-cyclodextrin (32 mg/ml)	93
	human serum albumin (40 mg/ml)	9
	QS-7 (250 µg/ml)	30
	aluminum hydroxide (2 mg/ml) in PBS	5
	aluminum hydroxide (2 mg/ml) in saline	13
20	Monophosphoryl lipid A (25 µg/ml)	4.9
	Monophosphoryl lipid A (50 µg/ml)	7.7
	Monophosphoryl lipid A (100 µg/ml)	6.5
	Triton X-100 (50 µg/ml)	1
	Triton X-100 (100 µg/ml)	1
25	Polysorbate 80 (2 mg/ml)	9
	Polysorbate 80 (4 mg/ml)	18
	Polysorbate 80 (10 mg/ml)	38

30

SUBSTITUTE SHEET (RULE 25)

Example 2**Compositions that Reduce the Lytic Effects of Other Saponins**

Other saponin adjuvants are also known to be hemolytic, although to
5 different extent than QS-21. These saponins include substantially pure QS-7,
QS-17, and QS-18. In addition, heterogeneous adjuvant saponins such as
Quil A are hemolytic. An example of the effect of Polysorbate 80 and
hydroxypropyl- β -cyclodextrin on the hemolytic indices of the substantially
10 pure QS-7 and heterogeneous Quil A is shown in Table 2. Hydroxypropyl- β -
cyclodextrin was shown to be effective in reducing the lytic effect (increasing
the hemolytic index) of QS-7. Polysorbate 80 and hydroxypropyl- β -
cyclodextrin were shown to be effective in minimizing the lytic effect
15 (increasing the hemolytic index) of Quil A.

Table 2:

Saponin	Excipient	Hemolytic Index ($\mu\text{g/ml}$)
20 QS-7	PBS	650
QS-7	Polysorbate 80 (8 mg/ml)	60
QS-7	Hydroxypropyl- β -cyclodextrin (32 mg/ml)	>1000
Quil A	PBS	18
Quil A	Polysorbate 80 (8 mg/ml)	43
25 Quil A	Hydroxypropyl- β -cyclodextrin (32 mg/ml)	200

Example 3

30

Compositions that Stabilize QS-21

QS-21 is an acylated bidesmodic triterpene saponin. It has a fatty acid
ester linked to the hydroxyl residues of fucose. In aqueous solution, this fatty

SUBSTITUTE SHEET (RULE 26)

acid ester migrates between two adjacent vicinal hydroxyl groups (fucose 3, 4) to form two equilibrium isomers (Jacobsen, N.E., Fairbrother, W.J., et al., 1996; Carbohydrate Research 280:1-14). The predominant isomer is acylated at fucose 4 and the minor isomer is acylated at fucose 3. This ester bond is the most labile bond in QS-21 and will hydrolyze under alkaline conditions to form a deacylated saponin and a fatty acid-arabinose domain. The deacylated saponin and the fatty acid domain are both inactive as immunological adjuvants (Kensil, C.R., et al., 1996, In: Saponins Used in Traditional and Modern Medicine, Waller and Yamaski, Eds., Plenum Press, NY, 165-172). Various conditions affect the stability of this ester bond (Cleland, J.L., et al., 1996, J. Pharmaceutical Sciences 85:22-28). Furthermore, the monomer form of QS-21 is more susceptible to hydrolysis than the micellar form.

Examples of the shelf life of QS-21 are shown in Table 3. The aqueous shelf life for a 50 µg/ml QS-21 solution at pH 7.0 at 4°C was shown to be only 94 days or approximately 3 months. This is representative of a typical clinical vaccine formulation containing QS-21 adjuvant (which consists of QS-21 at a concentration of 50-200 µg/ml in a physiological pH buffer (pH 7.0-7.5)). Hence, in simple buffer and salt solutions at low concentration, the QS-21 product does not maintain a desirable stability profile. Some improvement in stability, however, can be achieved by an increased concentration of the QS-21 product. For instance, the shelf life of a 500 µg/ml QS-21 solution at pH 7.0 at 4°C was shown to be 717 days, or 23.9 months. But a concentrated QS-21 solution is not necessarily a practical method of administering a low dose of adjuvant. For example, administration of 25 µg from a 500 µg/ml solution would require the syringe withdrawal of 0.05 ml of dose. Additionally, some

improved stability can be achieved by the use of a lower pH, i.e., at pH 6.0.

However, a pH substantially lower than the physiological pH range may not be tolerated well or be compatible with the antigen.

5 Table 3:

	QS-21 Concentration	pH	t_{90} (days)
	50 µg/ml	pH 7.0	94
	50 µg/ml	pH 6.0	679
10	500 µg/ml	pH 7.0	717

Another way to evaluate the stability of QS-21 in aqueous solution was to assay the solution by HPLC in an accelerated stability assay at 37°C.

15 Although this is not the temperature used for storage of vaccines (4°C), it was expected that this assay at 37°C would show the relative stabilizing power of a given excipient. For example, an excipient that extended the t_{90} value by two fold at 37°C would also be expected to extend the t_{90} value by two fold at 4°C.

20 Specifically, QS-21 (100 µg/ml) was prepared in various excipients in PBS at pH 7.0. The solutions were incubated at 37°C for 7 days. At the end of 7 days, the solutions were assayed by reversed phase-HPLC to determine the extent of degradation. The data was plotted as log (fraction QS-21 $t=7$ /QS-21 $t=0$ days) against time on the x-axis. The time to 10% degradation (t_{90}) was
25 extrapolated from this plot.

Table 4 shows the t_{90} values of QS-21 in various excipients. Stabilization of QS-21 is shown by an increase in t_{90} . Excipients that stabilized QS-21 by at
30 least two fold are Polysorbate 20, Polysorbate 80, native *Quillaja* saponin QS-7, and the deacylsaponin resulting from alkaline hydrolysis of QS-21 (DS-1).

Table 4:

	Excipient	t_{90} (days) at 37°C
	PBS (pH 7.0)	1.2
5	Polysorbate 20 (720 µg/ml)	2.9
	Polysorbate 80 (250 µg/ml)	3.2
	Polysorbate 80 (500 µg/ml)	4.3
	Polysorbate 80 (1.0 mg/ml)	5.2
	Polysorbate 80 (2.0 mg/ml)	7.2
10	Phenol (2.5 mg/ml)	2.3
	Pluronic F68 (1.0 mg/ml)	1.4
	QS-7 (100 µg/ml)	1.8
	QS-7 (250 µg/ml)	2.6
15	QS-7 (500 µg/ml)	9.0
	QS-7 (1.0 mg/ml)	16.0
	DS-1 (100 µg/ml)	2.2
	DS-1 (250 µg/ml)	3.3
	DS-1 (500 µg/ml)	7.2
20	DS-1 (1.0 mg/ml)	6.2
	Monocaproyl-rac-glycerol (1.0 mg/ml)	1.7
	α -cyclodextrin (5 mg/ml)	0.8
	β -cyclodextrin (5 mg/ml)	0.7
25	Methyl- β -cyclodextrin (5 mg/ml)	1.5
	hydroxypropyl- γ -cyclodextrin (5 mg/ml)	1.0
	hydroxypropyl- β -cyclodextrin (5 mg/ml)	1.0

In addition, 0.9% benzyl alcohol, and Plasdane C15 were evaluated for its ability to stabilize QS-21 (Table 5). All QS-21 concentrations and incubation conditions were equivalent in this experiment except that the QS-21 formulation was prepared in Dulbecco's PBS (without calcium or magnesium)

SUBSTITUTE SHEET (RULE 26)

at pH 7.5. As expected, the higher pH resulted in a faster degradation of QS-21 in PBS. However, Plasdane C15 stabilized QS-21.

5 Table 5:

	Excipient	t_{90} (days) at 37°C, pH 7.5
	Dulbecco's PBS	0.6
	0.9% benzyl alcohol in Dulbecco's PBS	0.7
10	Plasdane C15 in Dulbecco's PBS (25 mg/ml)	1.6
	Plasdane C15 in Dulbecco's PBS (50 mg/ml)	7.7

Example 4

15

Adjuvant Potency of Compositions

20

25

30

Figures 1A and 1B show the effect of Polysorbate 40, Polysorbate 60, Polysorbate 80, and methyl- β -cyclodextrin on the immune response of Balb/c mice to OVA plus various doses of QS-21. Female mice (10/group, 8-10 weeks of age at the first immunization) were immunized subcutaneously with 5 μ g of OVA and the indicated dose of QS-21 in either PBS alone or in 2 mg/ml excipient in PBS. A booster immunization was given by the same route at week 2. Sera was collected at week 4 for EIA analysis of the anti-OVA response. Mice were analyzed for OVA-specific IgG2a by a standard EIA analysis (Kensil, C.R., et al., 1993, Vaccine Research 2:273-281). QS-21 was active in all excipients within two fold of the threshold value determined in PBS. The same maximum level of antibody response was reached at the optimum adjuvant dose (typically 10 μ g and above).

SUBSTITUTE SHEET (RULE 26)

Figure 2 shows the effect of excipients on antibody response to a T-independent polysaccharide antigen. Balb/c mice were immunized subcutaneously with a commercial 23-valent *S. pneumonia* polysaccharide vaccine (Pnu-Imune, 0.5 µg/serotype) and different doses of QS-21 in PBS, in 4 mg/ml Polysorbate 80 in PBS, or in 16 mg/ml hydroxypropyl-β-cyclodextrin in PBS. Anti-Type 14 IgG was determined by EIA on sera collected at day 7 after a single immunization. Neither Polysorbate 80 or hydroxypropyl-β-cyclodextrin in the formulation reduced the potency of the vaccine for stimulating an IgG3 response specific for Type 14 polysaccharide serotype.

Example 5

Clinical Studies of Compositions-Trial 1

Various QS-21 compositions were administered to patients in order to test for the compositions' pain tolerance. Fifteen volunteers were recruited to receive four intramuscular injections, with each injection given at one week intervals. The study was carried out as a randomized, double-blind study. Three of the formulations contained 50 µg QS-21 in either Dulbecco's PBS (without calcium or magnesium), in 4 mg/ml Polysorbate 80 in PBS, or in 1 mg/ml aluminum hydroxide in saline. The fourth formulation was a PBS control without QS-21. Volunteers were asked to rate the immediate pain in the first five minutes after injection on a 0 to 10 scale (0=no pain, 1-3=mild, 4-7=moderate, 8-10=severe). The results are shown in Figure 3. The cumulative scores represented in Figure 3 of the patients' tolerance to pain is represented in Figure 4 as individual scores. The QS-21 formulation containing 4 mg/ml Polysorbate 80 resulted in an improved pain tolerance compared to QS-21 in

PBS. The highest score for this particular formulation was rated as a 5.

Example 6

5 Clinical Studies of Compositions-Trial 2

Various other QS-21 compositions were administered to patients in order to test for the compositions' pain tolerance. Fifteen volunteers were recruited to receive four intramuscular injections, with each injection given at
10 one week intervals. The study was carried out as a randomized, double-blind study. The excipients evaluated were benzyl alcohol, hydroxypropyl-beta-cyclodextrin, and a higher dose of Polysorbate 80, which had been shown to be
15 more effective than 4 mg/ml Polysorbate 80 at reducing QS-21 lysis of red blood cells *in vitro*. The five formulations tested were (1) 1 mg/ml aluminum hydroxide, which served as the placebo control; (2) 50 µg QS-21 in 0.72% benzyl alcohol in saline; (3) 50 µg QS-21 in 30 mg/ml hydroxypropyl-β-
20 cyclodextrin; (Encapsin, Janssen Biotech N.V., Olen, Belgium) (4) 50 µg QS-21 in 8 mg/ml Polysorbate 80; and (5) 50 µg QS-21 in PBS (Dulbecco's PBS without calcium or magnesium), which served as a positive control
25 formulation. Volunteers were asked to rate the immediate pain in the first five minutes after injection on a 0 to 10 scale (0=no pain, 1-3=mild, 4-7=moderate, 8-10=severe). The results are shown in Figure 5. The cumulative scores represented in Figure 5 of the patients' tolerance to pain is represented in
Figure 6 as individual scores. All excipients were shown to reduce the mean
30 and median pain scores associated with QS-21 in PBS. The highest single score for the QS-21/Encapsin formulation was rated as a 5, which compared more

favorably with the QS-21/Polysorbate 80 formulation that was rated with a single 6 and two 5's.

5 The invention now being fully described, it will be apparent to one of ordinary skill in the art that many changes and modifications can be made thereto without departing from the spirit or scope of the invention as set forth below.

10

15

20

25

30

We claim:

1. A composition comprising an antigen, a saponin adjuvant, and an excipient, wherein the composition reduces the *in vitro* lytic effect of the saponin adjuvant.
2. The composition according to claim 1, wherein the saponin adjuvant is a substantially pure saponin adjuvant.
3. The composition according to claim 2, wherein the substantially pure saponin adjuvant is QS-7 or QS-21.
4. The composition according to claim 1, wherein the saponin adjuvant is a heterogenous saponin adjuvant.
5. The composition according to claim 4, wherein the heterogeneous saponin adjuvant is Quil A.
6. The composition according to claim 1, wherein the antigen is a peptide, a protein, a polysaccharide, a lipid, or a nucleic acid encoding the peptide or protein.
7. The composition according to claim 1, wherein the excipient is a nonionic surfactant.
8. The composition according to claim 7, wherein the nonionic surfactant is a Polysorbate.
9. The composition according to claim 8, wherein the Polysorbate is Polysorbate 20, Polysorbate 40, Polysorbate 60, or Polysorbate 80.
10. The composition according to claim 1, wherein the excipient is a cyclodextrin.
11. The composition according to claim 10, wherein the cyclodextrin is

β -cyclodextrin.

12. The composition according to claim 11, wherein the β -cyclodextrin is hydroxypropyl- β -cyclodextrin.

5 13. The composition according to claim 1, wherein the composition further maintains the maximum adjuvant activity of QS-21.

14. The composition according to claim 13, wherein the saponin adjuvant is a substantially pure saponin adjuvant.

10 15. The composition according to claim 14, wherein the substantially pure saponin adjuvant is QS-7 or QS-21.

16. The composition according to claim 13, wherein the saponin adjuvant is a heterogenous saponin adjuvant.

15 17. The composition according to claim 16, wherein the heterogeneous saponin adjuvant is Quil A.

18. The composition according to claim 13, wherein the antigen is a
20 peptide, a protein, a polysaccharide, a lipid, or a nucleic acid encoding the peptide or protein.

19. The composition according to claim 13, wherein the excipient is a nonionic surfactant.

25 20. The composition according to claim 19, wherein the nonionic surfactant is a Polysorbate.

21. The composition according to claim 20, wherein the Polysorbate is Polysorbate 20, Polysorbate 40, Polysorbate 60, or Polysorbate 80.

30 22. The composition according to claim 13, wherein the excipient is a cyclodextrin.

23. The composition according to claim 22, wherein the cyclodextrin is β -cyclodextrin.

24. The composition according to claim 23, wherein the cyclodextrin is
5 hydroxypropyl- β -cyclodextrin.

25. The composition according to claim 1, wherein the composition further has an increased stability.

26. The composition according to claim 25, wherein the saponin
10 adjuvant is a substantially pure saponin adjuvant.

27. The composition according to claim 26, wherein the substantially pure saponin adjuvant is QS-7 or QS-21.

28. The composition according to claim 25, wherein the saponin
15 adjuvant is a heterogenous saponin adjuvant.

29. The composition according to claim 28, wherein the heterogeneous saponin adjuvant is Quil A.

20 30. The composition according to claim 25, wherein the antigen is a peptide, a protein, a polysaccharide, a lipid, or a nucleic acid encoding the peptide or protein.

31. The composition according to claim 25, wherein the excipient is a
25 nonionic surfactant.

32. The composition according to claim 31, wherein the nonionic surfactant is a Polysorbate.

33. The composition according to claim 26, wherein the Polysorbate is
30 Polysorbate 20, Polysorbate 40, Polysorbate 60, or Polysorbate 80.

34. The composition according to claim 1, wherein the composition

further improves the tolerance to saponin adjuvant associated pain in an individual to whom it is administered.

35. The composition according to claim 34, wherein the saponin
5 adjuvant is a substantially pure saponin.

36. The composition according to claim 35, wherein the substantially pure saponin adjuvant is QS-7 or QS-21.

37. The composition according to claim 34, wherein the saponin
10 adjuvant is a heterogenous saponin adjuvant.

38. The composition according to claim 37, wherein the heterogeneous saponin adjuvant is Quil A.

39. The composition according to claim 34, wherein the antigen is a
15 peptide, a protein, a polysaccharide, a lipid or a nucleic acid encoding the peptide or protein.

40. The composition according to claim 34, wherein the excipient is a
20 nonionic surfactant.

41. The composition according to claim 40, wherein the nonionic surfactant is a Polysorbate.

42. The composition according to claim 41, wherein the Polysorbate is
25 Polysorbate 20, Polysorbate 40, Polysorbate 60, or Polysorbate 80.

43. The composition according to claim 34, wherein the excipient is a cyclodextrin.

44. The composition according to claim 43, wherein the cyclodextrin is
30 β -cyclodextrin.

45. The composition according to claim 44, wherein the cyclodextrin is

hydroxypropyl- β -cyclodextrin.

5

10

15

20

25

30

1/7




FIG. 1A

RECTIFIED SHEET (RULE 91)
ISA/EP

2/7




FIG. 1B

RECTIFIED SHEET (RULE 91)
ISA/EP

3/7




FIG. 2

RECTIFIED SHEET (RULE 91)
ISA/EP

4/7




FIG. 3

RECTIFIED SHEET (RULE 91)
ISA/EP




FIG. 4

RECTIFIED SHEET (RULE 91)
ISA/EP

6/7




FIG. 5

RECTIFIED SHEET (RULE 91)
ISA/EP

7/7




FIG. 6

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 98/17940

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 A61K39/39 //(A61K39/39,47:40,47:34)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 650 398 A (SOLTYSIK SEAN ET AL) 22 July 1997 see column 1, line 12-17 see column 2, line 36-48 see column 6, line 10-24 see column 21, line 14-32 see column 22, line 3-36 see column 24, line 19-42	1-9, 13-21, 25-42
X	PATENT ABSTRACTS OF JAPAN vol. 095, no. 003, 28 April 1995 & JP 06 343419 A (DR MAINTSU:KK), 20 December 1994 see abstract -/-	1,2,4,6, 10

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" documents published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"A" document member of the same patent family

Date of the actual completion of the international search

20 January 1999

Date of mailing of the international search report

01/02/1999

Name and mailing address of the ISA
European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 551 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Covone, M

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 98/17940

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>WO 91 04052 A (PEPTIDE TECHNOLOGY LTD) 4 April 1991</p> <p>see page 3, line 19 - page 4, line 2 see page 4, line 28 - page 5, line 17 see claims 1-6</p>	<p>1,2,4-6, 25,26, 28-30, 34,35, 37-39</p>
X	<p>EP 0 187 286 A (BEHRINGWERKE AG) 16 July 1986</p> <p>see page 1, line 14 - page 2, line 4 see example 7 see claims 1,5,6</p>	<p>1-9, 13-21, 25-42</p>
X	<p>US 4 806 350 A (GERBER JAY D) 21 February 1989</p> <p>see column 1, line 64 - column 2, line 3 see column 3, line 47-59</p>	<p>1-9, 13-21, 25-42</p>
X	<p>GB 1 083 815 A (RALPH ARTHUR NEAL) 20 September 1967 see page 1, left-hand column, line 43 - right-hand column, line 52 see page 2, right-hand column, line 90-98</p>	<p>1,2,4-6</p>
A	<p>WO 96 20008 A (SOLVAY ;HILGERS LUUK (NL)) 4 July 1996</p> <p>see page 2, line 31-33 see page 7, line 5-33 see claims 1,2,4</p>	<p>10-12, 22-24, 43-45</p>
P,X	<p>WO 98 15287 A (SMITHKLINE BEECHAM BIOLOG ;FRIEDE MARTIN (BE); GARCON NATHALIE (BE) 16 April 1998</p> <p>see page 1, line 12-15 see page 2, line 3-7 see page 3, line 21-23 see example 5 see claims 1,2,15</p>	<p>1-6, 13-18, 25-30, 34-39</p>
E	<p>EP 0 884 056 A (SMITHKLINE BEECHAM BIOLOG) 16 December 1998</p> <p>see page 2, line 6-14 see page 4, line 24 - page 5, line 40</p>	<p>1-6, 13-18, 25-30, 34-39</p>

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/US 98/17940

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 5650398	A	22-07-1997	US 5443829 A	22-08-1995
			US 5273965 A	28-12-1993
			AU 6039396 A	18-12-1996
			CA 2223063 A	05-12-1996
			EP 0841932 A	20-05-1998
			WO 9638161 A	05-12-1996
			AU 687005 B	19-02-1998
			AU 4664993 A	31-01-1994
			CA 2139437 A	20-01-1994
			EP 0658111 A	21-06-1995
			FI 946173 A	15-02-1995
			HU 73814 A	30-09-1996
			JP 7509440 T	19-10-1995
			MX 9303977 A	29-04-1994
			NO 945071 A	01-03-1995
			WO 9401118 A	20-01-1995
WO 9104052	A	04-04-1991	AU 640414 B	26-08-1993
			AU 6421090 A	18-04-1991
			CA 2066653 A	23-03-1991
			EP 0491833 A	01-07-1992
EP 0187286	A	16-07-1986	DE 3446515 A	26-06-1986
			DK 593985 A	21-06-1986
			FI 855057 A	21-06-1986
			GR 853058 A	15-04-1986
			PT 81713 B	10-08-1987
US 4806350	A	21-02-1989	AU 609667 B	02-05-1991
			AU 7176087 A	22-10-1987
			CA 1282003 A	26-03-1991
			DE 3787663 D	11-11-1993
			DE 3787663 T	17-02-1994
			DK 190387 A	19-10-1987
			EP 0242205 A	21-10-1987
			ES 2059369 T	16-11-1994
			IE 61233 B	19-10-1994
			JP 2515321 B	10-07-1996
			JP 62255436 A	07-11-1987
			PT 84692 B	30-11-1989
GB 1083815	A		NONE	
WO 9620008	A	04-07-1996	BE 1008978 A	01-10-1996
			AU 4324896 A	19-07-1996
			AU 4324996 A	19-07-1996
			WO 9620222 A	04-07-1996
			BR 9510223 A	30-12-1997
			BR 9510263 A	04-11-1997
			CA 2208790 A	04-07-1996
			CA 2208849 A	04-07-1996
			CN 1171052 A	21-01-1998
			CN 1175264 A	04-03-1998
			EP 0814836 A	07-01-1998
			EP 0800539 A	15-10-1997
			JP 10511997 T	17-11-1998
WO 9815287	A	16-04-1998	AU 4781297 A	05-05-1998

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 98/17940

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0884056 A	16-12-1998	AU 693022 B	18-06-1998
		AU 5334596 A	18-11-1996
		AU 699213 B	26-11-1998
		AU 6987398 A	23-07-1998
		CA 2217178 A	31-10-1996
		CN 1182370 A	20-05-1998
		CZ 9703379 A	18-03-1998
		WO 9633739 A	31-10-1996
		EP 0822831 A	11-02-1998
		HU 9801560 A	28-10-1998
		NO 974859 A	21-10-1997
		PL 322968 A	02-03-1998
		SK 144297 A	06-05-1998